

THE MORPHOLOGY OF M17-UC1 – A DISK CANDIDATE SURROUNDING A HYPER-COMPACT H II REGION¹

M. NIELBOCK, R. CHINI, V. H. HOFFMEISTER, AND C. M. SCHEYDA
Astronomisches Institut, Ruhr-Universität Bochum, Universitätsstraße 150, 44780 Bochum, Germany

J. STEINACKER
Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany
Astronomisches Rechen-Institut am Zentrum für Astronomie Heidelberg, Mönchhofstraße 12-14, 69120 Heidelberg, Germany

D. NÜRNBERGER
European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile

R. SIEBENMORGEN
European Southern Observatory, Karl-Schwarzschild-Straße 2, 85748 Garching, Germany
Draft version February 5, 2008

ABSTRACT

We investigate the morphology and the evolutionary stage of the hyper-compact H II region M17-UC1 using observations at infrared wavelengths and NIR radiative transfer modelling. It is for the first time resolved into two emission areas separated by a dark lane reminiscent of an obscuring silhouette caused by a circumstellar disk. The observational data as well as model calculations so far suggest that M17-UC1 is surrounded by a disk of cool dust. This direct detection of a circumstellar disk candidate around a hyper-compact H II region is in agreement with the expectations of the disk accretion model for high-mass star formation.

Subject headings: stars: formation, circumstellar matter, pre-main sequence – open clusters and associations: general, individual (M17)

1. INTRODUCTION

1.1. General motivation

The hypothesis that high-mass stars form by accretion is supported by a number of indirect indicators like rotating molecular envelopes combined with bipolar outflows, gas in-fall or maser emission towards candidate high-mass protostellar objects (HMPOs) (e. g. Ho & Haschick 1986; Keto et al. 1987, 1988; Cesaroni et al. 1997, 2005; Zhang & Ho 1997; Zhang et al. 1998; Beltrán et al. 2004, 2005; Keto & Wood 2006). Although it seems plausible that such rotating and collapsing configurations will eventually form a disk, the involved observations usually lack the spatial resolution to firmly verify such a claim. Presently, there is no convincing example of a direct detection of a circumstellar disk around a HMPO that can be studied to infer the details of the accretion process including the ionisation effects. Even the currently most promising candidate, IRAS 20126+4104, must be questioned: Sridharan et al. (2005) report a bipolar structure, similar to what we present in this letter. However, Cesaroni et al. (2005) recently revised its mass to a value that lies just below what is commonly accepted as “high mass”. Furthermore, Hofner et al. (1999) explained their detection of radio continuum emission as shocked gas caused by a protostellar jet, although they also discussed free-free emission from ionised gas as a possible source.

In the present study, we investigated the hyper-compact H II region (HCHII) M17-UC1 (see Sect. 1.2 for details). HCHII are commonly associated with the earliest stages of high-mass star formation (Kurtz 2000; De Pree et al. 2005). They can be explained as a phase in early stellar evolution, when the high-mass star begins to ionise its molecular accretion flow. In such a stage, ionisation and accretion can co-exist (Keto 2003, 2006). We used near-IR observations in order to i) overcome the strong extinction towards the source, and ii) utilise adaptive optics techniques allowing for high spatial resolution. As a result, we find a bipolar structure reminiscent of a silhouette disk against a bright background as commonly found for low-mass protostars. Starting from this hypothesis, we perform radiative transfer modelling to verify this claim.

1.2. The object: M17-UC1

M17-UC1 was initially discovered as a cometary ultra-compact H II region by Felli et al. (1980, 1984), who suggested a B0 - B0.5 ZAMS star as the ionising source possessing a modelled Lyman continuum photon flux of $\simeq 2 \times 10^{47} \text{ s}^{-1}$. It was later re-classified as a HCHII with broad ($\geq 35 \text{ km s}^{-1}$) radio recombination lines and a rising spectral index of +1 between 1.4 and 43 GHz (Johnson et al. 1998; Sewilo et al. 2004). A number of class II methanol masers (Menten 1991) and hydroxyl masers (e.g. Churchwell et al. 1990; Caswell 1997; Walsh et al. 1998; Forster & Caswell 1999) are located in the vicinity of M17-UC1 being typical for the surroundings of high-mass protostars.

Within the positional accuracy of $5''.7$, Harper et al. (1976) detected a strong compact $10.6 \mu\text{m}$ source (IRC2) coinciding with M17-UC1 with an SED suggesting the presence of hot dust. The IR flux was assumed to originate entirely from

Electronic address: [nielbock, chini, vhoff, scheyda]@astro.rub.de
Electronic address: stein@mpia-hd.mpg.de
Electronic address: [dnuernbe, rsiebenm]@eso.org

¹ Based on observations made with ESO telescopes at the La Silla and Paranal observatories under programme IDs 71.C-0185(A), 71.C-0353(A), 73.C-0170(A), 73.C-0407(A), 75.C-0418(A), 77.C-0174(A)




FIG. 1.— M17-UC1 as seen by NACO (K_s and L') and VISIR (SiC/ N). The central position is 18:20:24.82, $-16:11:34.9$ (J2000), the field size is $10'' \times 10''$. The 3σ contour of the K_s band flux is superimposed on all images. The locations of associated OH ($\times$) and class II methanol masers ($+$) are indicated in the K_s band image. An enlargement ($1''.1 \times 1''.1$) of the source is shown in the upper right corner. The inset in the VISIR image illustrates the intensity distribution across the bright inner emission. Black contours at arbitrary intensity values are shown for enhancement. The lower right panel is a RGB coded composite of the three images (blue: K_s , green: L' , red: N).

M17-UC1 (Felli et al. 1984). When Felli & Stanga (1987) observed M17-UC1 from 1.25 to $18.1 \mu\text{m}$, they also found an unresolved IR source which they attributed to the HCHII. However, they used the strongest K band peak as their positional reference, which in fact is not M17-UC1 but the nearby IR-bright star IRS 5 (Chini & Wargau 1998). It is dominating the emission at JHK . Additional IR imaging was carried out by Giard et al. (1994), Chini et al. (2000) and Jiang et al. (2002). Kassis et al. (2002) characterised the source as a ZAMS B0 type surrounded by a shell of $0.6 - 3.4 M_\odot$ and a radius of less than 6600 AU.

2. OBSERVATIONS AND CALIBRATIONS

The $JHKsL'$ adaptive optics imaging was carried out in June 2003 using NAOS/CONICA (Lenzen et al. 2003; Rousset et al. 2003) at the ESO VLT. The pixel resolution was $0''.027$, the limiting magnitudes are $J = 20.6$, $H = 19.3$, $K_s = 18.4$ and $L' = 16.2$. The photometric calibration was obtained from our ISAAC observations in September 2002 (Chini et al. 2004; Hoffmeister et al. 2006). We adjusted the astrometry in the $JHKsL'$ images by referencing the NACO

sources with the detections of the ISAAC data that were astrometrically calibrated using the 2MASS database. With this procedure, we estimate a relative astrometric accuracy of better than $\pm 0''.1$.

The TIMMI2 (Reimann et al. 1998) NIR/MIR imaging was carried out during 3 observing runs in July 2003, April 2004, and July 2005 at the ESO 3.6 m telescope at La Silla, Chile. The observations covered the L , M , $N1$, $N10.4$, $N11.9$ and $Q1$ bands. A standard chopping and nodding technique was used with an amplitude of $10''$. All data are limited by diffraction with a FWHM of $0''.7$.

M17-UC1 was imaged with VISIR (Lagage et al. 2004) in May 2006 through the SiC filter. The pixel scale was $0''.127$ resulting in a field-of-view of $32''.5 \times 32''.5$. A standard chopping and nodding technique in perpendicular directions with throw amplitudes of $15''$ was applied. The measured image quality (FWHM) of $\simeq 0.32''$ is limited by diffraction. The astrometric calibration was achieved by fitting transformation equations between the NACO L' data and the VISIR data after identifying several point-sources visible in both wavebands. Image restoration of the TIMMI2 and VISIR data

was achieved with the software MOPSI (maintained by R. W. Zylka, IRAM, Grenoble, France). The overall astrometric accuracy is demonstrated by the RGB coded *KLN* composite image presented in Fig. 1.

The *K* band spectroscopy was obtained in August 2004 with ISAAC (Moorwood 1997) at a spectral resolution of 1500 and a slit width of $0''.3$. The spectroscopy of the *N* band silicate absorption feature was performed with TIMMI 2 at the ESO 3.6 m telescope at La Silla, Chile in July 2003. The seeing was $0''.7$; the slit width was $1''.2$.

3. RESULTS

3.1. Morphology

Fig. 1 displays the morphology of M17-UC1 at NIR and MIR wavelengths. On our images, M17-UC1 is barely visible at *H* with a brightness of 17.83 ± 0.25 mag. Surpassing previous NIR studies, we resolve the source for the first time into two *K* band emission blobs, separated by $0''.46$ at a position angle of 126° measured clockwise from north to south. A dark lane separates the two *K* band nebulae (Fig. 1). The south-western emission has an elliptical shape of $0''.9 \times 0''.5$ (3σ contour) and a pronounced peak at $18:20:24.83, -16:11:35.0$ (J2000) with a FWHM (full width at half maximum) of $0''.19 \times 0''.13$. The north-eastern emission at $18:20:24.85, -16:11:34.7$ (J2000) is more diffuse and has a size of $0''.8 \times 0''.5$. The integrated brightness of both blobs within a radius of 3600 AU is $K_s = 13.1$ mag, their intensity ratio is 10:1 (SW:NE). At *L* band, the source attains a spherical shape with a radius of $\sim 1''.0$ (3σ contour) and an *L'* brightness of 6.1 mag within 3600 AU.

On previous MIR images (Chini et al. 2000; Kassis et al. 2002), M17-UC1 appeared as a point-like source with spherically symmetric circumstellar emission. Our new *N* band data resolve M17-UC1 (see inset in Fig. 1). The peak brightness is centred on the south-western *K* band peak, and shows an elongation with a FWHM of $0''.7 \times 0''.5$ towards northeast being compatible with north-eastern *K* band peak. We derive a flux density of 29.87 ± 0.13 Jy for the central compact source. The total circumstellar emission ($\sim 5'' \times 3''$) is relatively complex with a noticeable extent to the northwest. It adds another 3.52 Jy to the total flux density.

3.2. Spectral appearance

Our *K* band spectrum (not shown) of M17-UC1 includes both nebulae and shows an extremely red continuum caused by hot dust. Apart from the $\text{Br}\gamma$ line emission, which most likely originates from scattered photons emitted by the gas inside the HCHII, the spectrum appears featureless.

In the *N* band, M17-UC1 displays a deep silicate feature (Fig. 2); its existence was already suggested by earlier photometric data (Harper et al. 1976; Kassis et al. 2002). This absorption indicates the presence of cool dust along the line of sight towards M17-UC1. Using the standard relation for converting optical depths of $\tau_\gamma \sim 17 \cdot \tau_{9.7}$ (Krügel 2003), we infer a visual extinction of about 40 mag. The luminosity between 1.6 and $20 \mu\text{m}$ corrected for extinction is $1.1 \times 10^4 L_\odot$ being consistent with an early B-type star. All our broad-band photometric data from previous epochs that were obtained with different filters trace the shape of the silicate absorption feature fairly well. This rules out possible MIR flux variations that were suggested by Nielbock et al. (2001). Although our broad *N* band imaging filter also includes the silicate feature, we only see the extended warm emitting dust in Fig. 1.




FIG. 2.— SED of M17-UC1. The data were obtained between 1999 and 2006. The red solid line is the MIR spectrum of the silicate absorption feature at $9.7 \mu\text{m}$. The inset is a zoom into the spectrum. A continuum fit of two black-body components with formal colour temperatures of $T = 425$ K and 225 K is shown as a dotted line.

4. DISCUSSION

4.1. Observational evidence for a disk

The symmetric absorption pattern of M17-UC1 in the *K* band along with the perpendicularly scattered light nebulosities on both sides very much resembles the appearance of young low-mass stars with circumstellar disks (e.g. McCaughrean & O'dell 1996; Padgett et al. 1999; Brandner et al. 2000; Grosso et al. 2003). This view is supported by the morphology of theoretically derived synthetic images of protostellar disks presented by Stark et al. (2006). These images are based on varying disk radius and accretion rate, among which one can find suiting representations of the *Ks* image of M17-UC1. In addition, the class II methanol masers that are located in the general direction of the supposed disk support our conjecture.

Apart from the almost uniformly extended *L* and *N* band emission one would expect to see the supposed disk to be emitting strongly in the MIR, too. Nevertheless, there must be cool dust inside the disk as witnessed by the silicate absorption feature. We interpret the elongated shape of the central MIR emission perpendicular to the disk orientation as scattered light from the central source. Due to the lower optical depth, however, the MIR disk emission is veiled by the silicate absorption and scattered MIR photons. At larger distances from the HCHII, we see a warmer and less dense envelope strongly emitting in the MIR.

Interestingly, stellar MIR emission is not necessarily attributed to disks, but can also originate from the walls of outflow cavities where the ambient interstellar medium interacts with the gas stream (De Buizer 2006, 2007). In fact, the *K* band nebulae could also be dominated by emission from an outflow with a wide opening angle. In this case, the narrow waist would indicate a small disk that restricts the outflow in these directions. However, our *K* band spectrum does not show any hints for shocks or outflows, so we discard this interpretation.

4.2. Model calculations

We have modelled the $2.2 \mu\text{m}$ radiation with a star-disk system using a 3D radiative transfer code described by Steinacker et al. (2006). The star was assumed to have $T_{\text{eff}} = 30000$ K. The disk has a radial power-law density profile with an exponential atmosphere above and below the disk mid-plane like




FIG. 3.— Model fit of M17-UC1. Left: Original Ks band image. Centre: Disk model inclined at 30° as discussed in the text. Right: Scattered light image at $2.2 \mu\text{m}$ from a filament in front of a B0 star. The filament has a diameter of about 3000 AU and a distance of 5000 AU to the star, the dust properties are identical to those of the disk model.

$$n(r, z) = n_0 \left(\frac{r}{r_0} \right)^\alpha \exp \left[- \left(\frac{z}{hr} \right)^2 \right] \quad (1)$$

with $r = \sqrt{x^2 + z^2}$, where n_0 is the number density normalisation, h the scale height, r_0 the inner radius, and r_1 the outer radius where the disk vanishes. The K band image with the triangular absorption is well represented by parameters $n_d = 10^4 \text{ m}^{-3}$, $\alpha = 1.3$, $h = 0.2$, $r_0 = 20 \text{ AU}$, $r_1 = 1000 \text{ AU}$ (Fig. 3) and standard $0.12 \mu\text{m}$ -sized silicate dust particles with opacities taken from Draine & Lee (1984).

The inclination of the disk is about 30° (edge-on = 0°). We obtain a lower mass limit of $4 \times 10^{-4} M_\odot$ from the scattered light that only probes the surface of the disk. A similar mass limit is derived from the extinction that blocks the stellar light at $2.2 \mu\text{m}$. Of course, more mass can be hidden in the interior of the disk without affecting the appearance of the object. Likewise, for a mildly edge-on disk, additional dust could reside outside the scattered light pattern.

We have also carried out radiative transfer calculations for the alternative scenario of a dusty filament that is unrelated to M17-UC1. We modelled dust configurations of differing sizes and masses located at various distances in front of the star. In contrast to the observed morphology, the scattered

light generally extends farther out along the filament, and the intensity ratio between the strong south-western peak and the faint north-eastern peak cannot be reproduced satisfactorily (Fig. 3). Therefore, the radiative transfer calculations also favour the disk model over the filament model.

5. CONCLUSION

We have presented new NIR and MIR observations of the hyper-compact HII region M17-UC1. As a prominent feature, the Ks band image shows a dark lane in scattered light producing a substantial silicate absorption feature observed in the MIR. Analysing the image with radiative transfer models, we find the best agreement by assuming a disk-like structure around the central source instead of a foreground filament. For this reason and because of the "high mass" nature of the embedded star as determined by many independent measurements, we suggest that M17-UC1 might be the first doubtless candidate of a HCHII where at least parts of a circumstellar disk are still present.

This work was partly funded by the Nordrhein-Westfälische Akademie der Wissenschaften. M. N. acknowledges the support by the Deutsche Forschungsgemeinschaft, project SFB 591 and the Ministerium für Innovation, Wissenschaft, Forschung und Technik (MIWFT) des Landes Nordrhein-Westfalen. We thank E. Churchwell, H. Beuther and M. Haas for helpful discussions. Likewise, we thank the referee E. Keto for his exemplary and valuable suggestions that significantly improved the manuscript. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

REFERENCES

- Beltrán, M. T., Cesaroni, R., Neri, R., et al. 2004, *ApJ*, 601, L187
 Beltrán, M. T., Cesaroni, R., Neri, R., et al. 2005, *A&A*, 435, 901
 Brandner, W., Sheppard, S., Zinnecker, H., et al. 2000, *A&A*, 364, L13
 Caswell, J. L. 1997, *MNRAS*, 289, 203
 Cesaroni, R., Felli, M., Testi, L., Walmsley, C. M., & Olmi, L. 1997, *A&A*, 325, 725
 Cesaroni, R., Neri, R., Olmi, L., et al. 2005, *A&A*, 434, 1039
 Chini, R., Hoffmeister, V. H., Kämpgen, K., et al. 2004, *A&A*, 427, 849
 Chini, R., Nielbock, M., & Beck, R. 2000, *A&A*, 357, L33
 Chini, R. & Wargau, W. F. 1998, *A&A*, 329, 161
 Churchwell, E., Walmsley, C. M., & Cesaroni, R. 1990, *A&AS*, 83, 119
 De Buizer, J. M. 2006, *ApJ*, 642, L57
 De Buizer, J. M. 2007, *ApJ*, 654, L147
 De Pree, C. G., Wilner, D. J., Deblasio, J., Mercer, A. J., & Davis, L. E. 2005, *ApJ*, 624, L101
 Draine, B. T. & Lee, H. M. 1984, *ApJ*, 285, 89
 Felli, M., Johnston, K. J., & Churchwell, E. 1980, *ApJ*, 242, L157
 Felli, M., Massi, M., & Churchwell, E. 1984, *A&A*, 136, 53
 Felli, M. & Stanga, R. 1987, *A&A*, 175, 193
 Forster, J. R. & Caswell, J. L. 1999, *A&AS*, 137, 43
 Giard, M., Bernard, J. P., Lacombe, F., Normand, P., & Rouan, D. 1994, *A&A*, 291, 239
 Grosso, N., Alves, J., Wood, K., et al. 2003, *ApJ*, 586, 296
 Harper, D. A., Low, F. J., Rieke, G. H., & Thronson, H. A. 1976, *ApJ*, 205, 136
 Ho, P. T. P. & Haschick, A. D. 1986, *ApJ*, 304, 501
 Hoffmeister, V. H., Chini, R., Scheyda, C. M., et al. 2006, *A&A*, 457, L29
 Hofner, P., Cesaroni, R., Rodríguez, L. F., & Martí, J. 1999, *A&A*, 345, L43
 Jiang, Z., Yao, Y., Yang, J., et al. 2002, *ApJ*, 577, 245
 Johnson, C. O., De Pree, C. G., & Goss, W. M. 1998, *ApJ*, 500, 302
 Kassis, M., Deutsch, L. K., Campbell, M. F., et al. 2002, *AJ*, 124, 1636
 Keto, E. 2003, *ApJ*, 599, 1196
 Keto, E. 2006, astro-ph/0603856
 Keto, E. & Wood, K. 2006, *ApJ*, 637, 850
 Keto, E. R., Ho, P. T. P., & Haschick, A. D. 1988, *ApJ*, 324, 920
 Keto, E. R., Ho, P. T. P., & Reid, M. J. 1987, *ApJ*, 323, L117
 Krügel, E. 2003, *The Physics of Interstellar Dust*, Series in Astronomy and Astrophysics (Bristol, UK: Institute of Physics Publishing)
 Kurtz, S. E. 2000, in *Revista Mexicana de Astronomía y Astrofísica (Serie de Conferencias)*, Vol. 9, Astrophysical Plasmas: Codes, Models, and Observations, ed. J. Arthur, N. Brickhouse, & J. Franco (Instituto de Astronomía, UNAM, Mexico), 169–176
 Lagage, P. O., Pel, J. W., Authier, M., et al. 2004, *The Messenger*, 117, 12
 Lenzen, R., Hartung, M., Brandner, W., et al. 2003, in *Proceedings of the SPIE*, Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes, ed. M. Iye & A. M. F. Moorwood (The International Society for Optical Engineering), 944–952
 McCaughrean, M. J. & O'dell, C. R. 1996, *AJ*, 111, 1977
 Menten, K. M. 1991, *ApJ*, 380, L75
 Moorwood, A. F. 1997, in *Proceedings of the SPIE*, Vol. 2871, Optical Telescopes of Today and Tomorrow, ed. A. L. Ardeberg (The International Society for Optical Engineering), 1146–1151
 Nielbock, M., Chini, R., Jütte, M., & Manthey, E. 2001, *A&A*, 377, 273
 Padgett, D. L., Brandner, W., Stapelfeldt, K. R., et al. 1999, *AJ*, 117, 1490
 Reimann, H., Weinert, U., & Wagner, S. 1998, in *Proceedings of the SPIE*, Vol. 3354, Infrared Astronomical Instrumentation, ed. A. M. Fowler (The International Society for Optical Engineering), 865–876
 Rousset, G., Lacombe, F., Puget, P., et al. 2003, in *Proceedings of the SPIE*, Vol. 4839, Adaptive Optical System Technologies II, ed. P. L. Wizinowich & D. Bonaccini (The International Society for Optical Engineering), 140–149
 Sewilo, M., Churchwell, E., Kurtz, S., Goss, W. M., & Hofner, P. 2004, *ApJ*, 605, 285
 Sridharan, T. K., Williams, S. J., & Fuller, G. A. 2005, *ApJ*, 631, L73

- Stark, D. P., Whitney, B. A., Stassun, K., & Wood, K. 2006, *ApJ*, 649, 900
 Steinacker, J., Chini, R., Nielbock, M., et al. 2006, *A&A*, 456, 1013
 Walsh, A. J., Burton, M. G., Hyland, A. R., & Robinson, G. 1998, *MNRAS*, 301, 640
 Zhang, Q. & Ho, P. T. P. 1997, *ApJ*, 488, 241
 Zhang, Q., Hunter, T. R., & Sridharan, T. K. 1998, *ApJ*, 505, L151